

Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

# Custom Soil Resource Report for Robeson County, North Carolina

**Cowpen Swamp Road Farm** 



# **Preface**

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2 053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

# **Contents**

Preface	2
How Soil Surveys Are Made	
Soil Map	
Soil Map	
Legend	
Map Unit Legend	
Map Unit Descriptions	
Robeson County, North Carolina	13
BB—Bibb soils	
GoA—Goldsboro loamy sand, 0 to 2 percent slopes, Southern Coastal	
Plain	14
Ly—Lynchburg sandy loam, 0 to 2 percent slopes	15
NoA—Norfolk loamy sand, 0 to 2 percent slopes	17
NoB—Norfolk loamy sand, 2 to 6 percent slopes	18
Ra—Rains sandy loam, 0 to 2 percent slopes	19
W—Water	21
WaB—Wagram loamy sand, 0 to 6 percent slopes	22
Soil Information for All Uses	24
Suitabilities and Limitations for Use	24
Vegetative Productivity	24
Yields of Non-Irrigated Crops (Component): Corn (Bu)	24
Yields of Non-Irrigated Crops (Component): Soybeans (Bu)	28
References	33

# **How Soil Surveys Are Made**

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

# Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



# MAP LEGEND

# Area of Interest (AOI)

Area of Interest (AOI)

#### Soils

Soil Map Unit Polygons

Soil Map Unit Lines

Soil Map Unit Points

#### **Special Point Features**

ဖ

Blowout

Borrow Pit

Clay Spot

**Closed Depression** 

Gravel Pit

**Gravelly Spot** 

Landfill Lava Flow

Marsh or swamp

Mine or Quarry

Miscellaneous Water Perennial Water

Rock Outcrop

Saline Spot

Sandy Spot

Severely Eroded Spot

Sinkhole

Sodic Spot

Slide or Slip

Spoil Area



Very Stony Spot

Stony Spot



Wet Spot Other



Special Line Features

# Water Features

Streams and Canals

# Transportation

---

Rails

Interstate Highways

**US Routes** 

Major Roads

00

Local Roads

# Background

Aerial Photography

# MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Robeson County, North Carolina Survey Area Data: Version 22, Sep 13, 2023

Soil map units are labeled (as space allows) for map scales 1:50.000 or larger.

Date(s) aerial images were photographed: Apr 17, 2022—May 20. 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

# **Map Unit Legend**

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI		
ВВ	Bibb soils	11.5	8.9%		
GoA	Goldsboro loamy sand, 0 to 2 percent slopes, Southern Coastal Plain	22.2	17.2%		
Ly	Lynchburg sandy loam, 0 to 2 percent slopes	5.9	4.6%		
NoA	Norfolk loamy sand, 0 to 2 percent slopes	53.2	41.1%		
NoB	Norfolk loamy sand, 2 to 6 percent slopes	0.3	0.2%		
Ra	Rains sandy loam, 0 to 2 percent slopes	21.7	16.8%		
W	Water	0.2	0.2%		
WaB	Wagram loamy sand, 0 to 6 percent slopes	14.2	11.0%		
Totals for Area of Interest		129.2	100.0%		

# **Map Unit Descriptions**

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a

given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An association is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

# **Robeson County, North Carolina**

# **BB**—Bibb soils

# **Map Unit Setting**

National map unit symbol: 3vdw

Elevation: 80 to 330 feet

Mean annual precipitation: 38 to 55 inches Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 210 to 265 days

Farmland classification: Not prime farmland

# **Map Unit Composition**

Bibb, undrained, and similar soils: 80 percent Johnston, undrained, and similar soils: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Bibb, Undrained**

# Setting

Landform: Flood plains

Landform position (two-dimensional): Toeslope

Down-slope shape: Concave Across-slope shape: Linear

Parent material: Sandy and loamy alluvium

# **Typical profile**

A - 0 to 6 inches: sandy loam

Cg1 - 6 to 60 inches: sandy loam

Cg2 - 60 to 80 inches: loamy sand

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Frequent Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 7.2 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5w

Hydrologic Soil Group: A/D Hydric soil rating: Yes

# **Description of Johnston, Undrained**

# Setting

Landform: Flood plains
Down-slope shape: Concave
Across-slope shape: Linear

Parent material: Sandy and loamy alluvium

# Typical profile

A - 0 to 30 inches: mucky loam

Cg1 - 30 to 34 inches: loamy fine sand Cg2 - 34 to 80 inches: fine sandy loam

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Ponded

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: About 0 inches Frequency of flooding: Frequent Frequency of ponding: Frequent

Available water supply, 0 to 60 inches: High (about 9.4 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7w

Hydrologic Soil Group: A/D Hydric soil rating: Yes

# GoA—Goldsboro loamy sand, 0 to 2 percent slopes, Southern Coastal Plain

# Map Unit Setting

National map unit symbol: 2v750

Elevation: 110 to 300 feet

Mean annual precipitation: 40 to 55 inches
Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: All areas are prime farmland

# **Map Unit Composition**

Goldsboro and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Goldsboro**

# **Setting**

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Loamy marine deposits

# Typical profile

Ap - 0 to 9 inches: loamy sand

E - 9 to 12 inches: loamy sand

Bt - 12 to 62 inches: sandy clay loam

Btg - 62 to 80 inches: sandy clay loam

# Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.57 to 1.98 in/hr)

Depth to water table: About 24 to 36 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 8.1 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: B

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# **Minor Components**

# Norfolk

Percent of map unit: 8 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex, linear Across-slope shape: Convex, linear

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# Lynchburg

Percent of map unit: 7 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# Ly—Lynchburg sandy loam, 0 to 2 percent slopes

# Map Unit Setting

National map unit symbol: 2vx8p

Elevation: 10 to 330 feet

Mean annual precipitation: 40 to 55 inches
Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: Prime farmland if drained

# **Map Unit Composition**

Lynchburg and similar soils: 84 percent

Minor components: 16 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Lynchburg**

# Setting

Landform: Flats on marine terraces, broad interstream divides on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Loamy marine deposits

# Typical profile

Ap - 0 to 8 inches: sandy loam
E - 8 to 11 inches: sandy loam
Bt - 11 to 21 inches: sandy clay loam
Btg - 21 to 65 inches: sandy clay loam
BCg - 65 to 85 inches: sandy clay loam

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.57 to 1.98 in/hr)

Depth to water table: About 6 to 18 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 6.6 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: B/D

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# **Minor Components**

## Rains

Percent of map unit: 8 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Dip, talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

# Goldsboro

Percent of map unit: 8 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# NoA—Norfolk loamy sand, 0 to 2 percent slopes

# **Map Unit Setting**

National map unit symbol: 2v75w

Elevation: 10 to 330 feet

Mean annual precipitation: 40 to 55 inches Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: All areas are prime farmland

# **Map Unit Composition**

Norfolk and similar soils: 83 percent Minor components: 17 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Norfolk**

# Setting

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex, linear Across-slope shape: Convex, linear Parent material: Loamy marine deposits

# Typical profile

Ap - 0 to 8 inches: loamy sand E - 8 to 14 inches: loamy sand Bt - 14 to 65 inches: sandy clay loam BC - 65 to 80 inches: sandy clay loam

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.57 to 1.98 in/hr)

Depth to water table: About 40 to 72 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 6.9 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 1

Hydrologic Soil Group: A

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# **Minor Components**

#### Goldsboro

Percent of map unit: 9 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# Wagram

Percent of map unit: 8 percent

Landform: Ridges on marine terraces, broad interstream divides on marine

terraces

Landform position (two-dimensional): Summit, shoulder Landform position (three-dimensional): Crest, talf

Down-slope shape: Convex Across-slope shape: Convex

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# NoB—Norfolk loamy sand, 2 to 6 percent slopes

# Map Unit Setting

National map unit symbol: 2v75y

Elevation: 30 to 450 feet

Mean annual precipitation: 38 to 55 inches Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: All areas are prime farmland

# **Map Unit Composition**

Norfolk and similar soils: 83 percent Minor components: 17 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Norfolk**

# Settina

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex, linear Across-slope shape: Convex, linear Parent material: Loamy marine deposits

# **Typical profile**

Ap - 0 to 8 inches: loamy sand E - 8 to 14 inches: loamy sand Bt - 14 to 65 inches: sandy clay loam

BC - 65 to 80 inches: sandy clay loam

# **Properties and qualities**

Slope: 2 to 6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.57 to 1.98 in/hr)

Depth to water table: About 40 to 72 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 6.9 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: A

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# **Minor Components**

# Wagram

Percent of map unit: 10 percent

Landform: Ridges on marine terraces, broad interstream divides on marine

terraces

Landform position (two-dimensional): Summit, shoulder Landform position (three-dimensional): Riser, rise

Down-slope shape: Convex, linear Across-slope shape: Convex

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# Goldsboro

Percent of map unit: 7 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# Ra—Rains sandy loam, 0 to 2 percent slopes

# **Map Unit Setting**

National map unit symbol: 2v760

Elevation: 30 to 330 feet

Mean annual precipitation: 40 to 55 inches Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: Prime farmland if drained

# **Map Unit Composition**

Rains, undrained, and similar soils: 58 percent Rains, drained, and similar soils: 24 percent

Minor components: 18 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Rains, Undrained**

# Setting

Landform: Broad interstream divides on marine terraces, flats on marine terraces,

carolina bays on marine terraces

Landform position (three-dimensional): Dip, talf

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Loamy marine deposits

# **Typical profile**

A - 0 to 6 inches: sandy loam

Eg - 6 to 12 inches: sandy loam

Btg - 12 to 65 inches: sandy clay loam

BCg - 65 to 80 inches: sandy clay loam

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 1.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 7.9 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: A/D

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

# **Description of Rains, Drained**

## Settina

Landform: Broad interstream divides on marine terraces, flats on marine terraces, carolina bays on marine terraces

Landform position (three-dimensional): Dip, talf

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Loamy marine deposits

# Typical profile

Ap - 0 to 6 inches: sandy loam

Eg - 6 to 12 inches: sandy loam

Btg - 12 to 65 inches: sandy clay loam

BCg - 65 to 80 inches: sandy clay loam

# **Properties and qualities**

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 1.98 in/hr)

Depth to water table: About 12 to 36 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 7.9 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3w

Hydrologic Soil Group: B

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

# **Minor Components**

# Lynchburg

Percent of map unit: 10 percent

Landform: Broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear Across-slope shape: Linear

Ecological site: F153AY040NC - Moist Loamy Rises and Flats

Hydric soil rating: No

# Pantego, undrained

Percent of map unit: 8 percent

Landform: Flats, broad interstream divides, stream terraces

Landform position (three-dimensional): Tread, talf

Down-slope shape: Linear Across-slope shape: Concave

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

# W-Water

# **Map Unit Composition**

Water: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Water**

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 8

Hydric soil rating: No

# WaB—Wagram loamy sand, 0 to 6 percent slopes

# **Map Unit Setting**

National map unit symbol: 3vg3

Elevation: 80 to 330 feet

Mean annual precipitation: 38 to 55 inches
Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 210 to 265 days

Farmland classification: Farmland of statewide importance

# **Map Unit Composition**

Wagram and similar soils: 90 percent

Minor components: 5 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

# **Description of Wagram**

# Setting

Landform: Ridges on marine terraces, broad interstream divides on marine

terraces

Landform position (two-dimensional): Summit, shoulder

Landform position (three-dimensional): Crest

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Loamy marine deposits

# **Typical profile**

Ap - 0 to 8 inches: loamy sand E - 8 to 24 inches: loamy sand Bt - 24 to 75 inches: sandy clay loam BC - 75 to 83 inches: sandy loam

# **Properties and qualities**

Slope: 0 to 6 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.57 to 1.98 in/hr)

Depth to water table: About 60 to 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 6.7 inches)

# Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2s

Hydrologic Soil Group: A

Ecological site: F153AY030NC - Dry Loamy Rises and Flats

Hydric soil rating: No

# **Minor Components**

# Bibb, undrained

Percent of map unit: 3 percent

Landform: Flood plains

Landform position (two-dimensional): Toeslope

Down-slope shape: Concave Across-slope shape: Linear

Ecological site: F153AY090NC - Flooded Mineral Soil Floodplains and Terraces

Hydric soil rating: Yes

# Johnston, undrained

Percent of map unit: 2 percent Landform: Flood plains Down-slope shape: Concave Across-slope shape: Linear

Ecological site: F153AY090NC - Flooded Mineral Soil Floodplains and Terraces

Hydric soil rating: Yes

# Soil Information for All Uses

# Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

# Vegetative Productivity

Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

# Yields of Non-Irrigated Crops (Component): Corn (Bu)

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.


The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative"

value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.



# MAP LEGEND

# Area of Interest (AOI)

Area of Interest (AOI)

#### Soils

# Soil Rating Polygons

<= 45.00

> 45.00 and <= 70.11

> 70.11 and <= 111.95

> 111.95 and <= 113.64

Not rated or not available

> 113.64 and <= 128.45

#### Soil Rating Lines

<= 45.00

> 45.00 and <= 70.11

> 70.11 and <= 111.95

> 111.95 and <= 113.64

> 113.64 and <= 128.45

Not rated or not available

## **Soil Rating Points**

<= 45.00

> 45.00 and <= 70.11

> 70.11 and <= 111.95

> 111.95 and <= 113.64

> 113.64 and <= 128.45

Not rated or not available

#### Water Features

Streams and Canals

# Transportation

Interstate Highways

US Routes

Major Roads

Local Roads

#### Background

Aerial Photography

# MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service

Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Robeson County, North Carolina Survey Area Data: Version 22, Sep 13, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 17, 2022—May 20, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

# Table—Yields of Non-Irrigated Crops (Component): Corn (Bu)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
ВВ	Bibb soils		11.5	8.9%
GoA	Goldsboro loamy sand, 0 to 2 percent slopes, Southern Coastal Plain	128.45	22.2	17.2%
Ly	Lynchburg sandy loam, 0 to 2 percent slopes	113.64	5.9	4.6%
NoA	Norfolk loamy sand, 0 to 2 percent slopes	113.07	53.2	41.1%
NoB	Norfolk loamy sand, 2 to 6 percent slopes	111.95	0.3	0.2%
Ra	Rains sandy loam, 0 to 2 percent slopes	45.00	21.7	16.8%
W	Water		0.2	0.2%
WaB	Wagram loamy sand, 0 to 6 percent slopes	70.11	14.2	11.0%
Totals for Area of Interest			129.2	100.0%

# Rating Options—Yields of Non-Irrigated Crops (Component): Corn (Bu)

Crop: Corn Yield Units: Bu

Aggregation Method: Weighted Average
Component Percent Cutoff: None Specified

Tie-break Rule: Higher
Interpret Nulls as Zero: Yes

# Yields of Non-Irrigated Crops (Component): Soybeans (Bu)

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.


In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.

The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.



# MAP LEGEND

# Area of Interest (AOI)

Area of Interest (AOI)

## Soils

# Soil Rating Polygons

<= 15.16

> 15.16 and <= 25.58

> 25.58 and <= 41.07

> 41.07 and <= 42.07

Not rated or not available

> 42.07 and <= 44.83

#### Soil Rating Lines

<= 15.16

> 15.16 and <= 25.58

> 25.58 and <= 41.07

> 41.07 and <= 42.07

> 42.07 and <= 44.83

Not rated or not available

## **Soil Rating Points**

<= 15.16

> 15.16 and <= 25.58

> 25.58 and <= 41.07

> 41.07 and <= 42.07

> 42.07 and <= 44.83

Not rated or not available

#### **Water Features**

Streams and Canals

# Transportation

→ Rails

. .

Interstate Highways

US Routes

Major Roads

Local Roads

#### Background

Aerial Photography

# MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service

Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Robeson County, North Carolina Survey Area Data: Version 22, Sep 13, 2023

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 17, 2022—May 20, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

# Table—Yields of Non-Irrigated Crops (Component): Soybeans (Bu)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
ВВ	Bibb soils		11.5	8.9%
GoA	Goldsboro loamy sand, 0 to 2 percent slopes, Southern Coastal Plain	44.83	22.2	17.2%
Ly	Lynchburg sandy loam, 0 to 2 percent slopes	42.07	5.9	4.6%
NoA	Norfolk loamy sand, 0 to 2 percent slopes	41.07	53.2	41.1%
NoB	Norfolk loamy sand, 2 to 6 percent slopes	40.71	0.3	0.2%
Ra	Rains sandy loam, 0 to 2 percent slopes	15.16	21.7	16.8%
W	Water		0.2	0.2%
WaB	Wagram loamy sand, 0 to 6 percent slopes	25.58	14.2	11.0%
Totals for Area of Interest			129.2	100.0%

# Rating Options—Yields of Non-Irrigated Crops (Component): Soybeans (Bu)

Crop: Soybeans
Yield Units: Bu

Aggregation Method: Weighted Average Component Percent Cutoff: None Specified

Tie-break Rule: Higher Interpret Nulls as Zero: Yes

# References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2\_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcs142p2\_052290.pdf